

The earliest segmental sternum in a Permian synapsid and its implications for the evolution of mammalian locomotion and ventilation

Eva-Maria Bendel^{*1,2}, Christian F. Kammerer^{3,4}, Zhe-Xi Luo⁵, Roger M.H. Smith^{4,6}, Jörg Fröbisch^{1,2,4}

¹Museum für Naturkunde, Leibniz-Institut für Evolutions- und Biodiversitätsforschung, Invalidenstraße 43, 10115 Berlin, Germany

²Institut für Biologie, Humboldt-Universität zu Berlin, Invalidenstraße 42, 10115 Berlin, Germany

³North Carolina Museum of Natural Sciences, 11 W Jones Street, Raleigh, NC, USA

⁴Evolutionary Studies Institute, University of the Witwatersrand, Yale Road, Johannesburg, 2000, South Africa

⁵Department of Organismal Biology and Anatomy, University of Chicago, 1027 E 57th Street, Chicago, IL, USA

⁶Department of Karoo Palaeontology, Iziko South African Museum, 25 Queen Victoria Street, Cape Town, 8001, South Africa


* Eva-Maria Bendel.

Supplementary material.

Supplementary Table 1.


Basic measurements of skull of *Gorgonops torvus* (specimen SAM-PK K10591)

Skull length (estimated):	ca. 30 cm
Snout length:	ca. 13 cm
Height of snout:	ca. 10 cm (measured from ventral most part of maxilla to skull roof)
Skull width:	ca. 6.5 cm (although extremely laterally compressed)



a: *Gorgonops torvus*
SAM-PK-K10591, dorsal view

b: "Aelurognathus" microdon
SAM-PK-9344, ventral view



c: *Cyonosaurus* sp.
USNM 412381, ventral view

d: undescribed gorgonopsian
NHCC LB350, ventral view

Supplementary Figure 1. Gorgonopsians preserving the sternum and/or interclavicle, used as a base for the reconstruction in Fig. 2b. Photograph in SFig 1d adapted from Christian A. Sidor, all rights reserved.